



## King's Research Portal

DOI:

[10.1016/j.schres.2018.06.008](https://doi.org/10.1016/j.schres.2018.06.008)

*Document Version*

Peer reviewed version

[Link to publication record in King's Research Portal](#)

*Citation for published version (APA):*

Taylor, M. J., & Yim, S. (2018). Is there rebound psychosis on withdrawal of antipsychotic medication in schizophrenia? *Schizophrenia Research*. <https://doi.org/10.1016/j.schres.2018.06.008>

### **Citing this paper**

Please note that where the full-text provided on King's Research Portal is the Author Accepted Manuscript or Post-Print version this may differ from the final Published version. If citing, it is advised that you check and use the publisher's definitive version for pagination, volume/issue, and date of publication details. And where the final published version is provided on the Research Portal, if citing you are again advised to check the publisher's website for any subsequent corrections.

### **General rights**

Copyright and moral rights for the publications made accessible in the Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognize and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the Research Portal

### **Take down policy**

If you believe that this document breaches copyright please contact [librarypure@kcl.ac.uk](mailto:librarypure@kcl.ac.uk) providing details, and we will remove access to the work immediately and investigate your claim.

## Letter to the editor

Is there rebound psychosis on withdrawal of antipsychotic medication in schizophrenia?

Matthew J Taylor MRCPsych <sup>1,2</sup>

Seb Yim <sup>1</sup>

1. Institute of Psychiatry, Psychology & Neuroscience, King's College London, London SE5 8AF, UK
2. Psychosis Clinical Academic Group, South London and Maudsley NHS Foundation Trust, London SE5 8AZ, UK

Author to whom correspondence should be addressed:

Dr Taylor, Department of Psychosis Studies, IOPPN PO63, De Crespigny Park, King's College London, London SE5 8AF, United Kingdom

Tel. +44 20 7848 0363

Fax. +44 20 7848 0976

email: [matthew.j.taylor@kcl.ac.uk](mailto:matthew.j.taylor@kcl.ac.uk)

## Letter

Dear Editors:

It is well established that dopamine antagonist medications, antipsychotics, have an important role in the prevention of relapse in schizophrenia and other psychoses (Buchanan et al., 2010; Hasan et al., 2013). Meta-analysis of randomised controlled trials indicates continuation of medication is associated with fewer relapses over the next year than discontinuation (Leucht et al., 2012). However, there remains vigorous ongoing debate around longer-term risks and benefits (Correll et al., 2018), and whether adaptive changes such as dopamine supersensitivity may lead to rebound psychosis on medication withdrawal, which could inflate the apparent benefits of continuing medication (Murray et al., 2016).

Some previous reviews including observational studies have suggested relapse rates are most substantially elevated in the first three months following medication discontinuation, consistent with an early rebound effect. (Baldessarini et al., 1995), analogous with that previously described for lithium withdrawal in bipolar disorder (Suppes et al., 1991). We recently analysed data from randomised trials comparing continuation and discontinuation strategies in people with schizophrenia, using time to event data to allow comparison of effects between different time periods post-randomisation.

Clinical trials in which people with a diagnosis of schizophrenia stable on medication were randomised to either continue on, or withdraw, an antipsychotic medication, were identified from a systematic review reported previously (Leucht et al., 2012). Of these studies the subset presenting time to event data on relapse as survival curves were identified. Survival curve data were extracted using a plot digitizer

(<https://automeris.io/WebPlotDigitizer>, version 4.1). Interval Hazard Ratios (HR) and their variance were calculated as previously described (Parmar et al., 1998; Tierney et al., 2007). From studies providing data over the first six months following randomisation, pooled estimates of HR and 95% Confidence Intervals (CI) were obtained by inverse variance meta-analysis with a random effects model. Heterogeneity was assessed by use of  $I^2$ . Analyses were performed using *R* (version 3.5), with *meta* package (version 4.9).

12 randomised controlled trials, in which 2656 participants with schizophrenia had been randomised to continue or discontinue medication, provided survival curve data suitable for analysis over the six month period following randomisation.


Figure 1 near here

People continuing antipsychotic medication were at lower risk of relapse over the six months following randomization - HR 0.38 (95% CI 0.32 to 0.44). A similar effect was seen across times studied (Figure 1). There was no significant effect of time on HR ( $p=.3$ ). Within the first month after randomisation, HR 0.46 (95% CI 0.36 to 0.59).

These data from meta-analysis of randomised controlled trials confirm a lower risk of relapse in those randomised to continue antipsychotic medications. HRs within the first six months are similar to the overall estimates of risk ratio (0.40, 95% CI 0.33 to 0.49) previously described (Leucht et al., 2012). Contrary to previous reports (Baldessarini et al., 1995), no evidence of excess early relapse in those stopping medication was seen - the consistent hazard ratio over time indicates no general early rebound phenomenon is seen, although a specific effect of particular agents, or in particular populations, cannot be excluded by this overall analysis. Future studies could specifically investigate those treated with clozapine, treatment-resistant

schizophrenia, where apparent rebound is most often suspected (Chouinard et al., 2017).

Hazard Ratio vs placebo



## Legends

### Figure 1

Risk of relapse (Hazard ratios with 95% Confidence Intervals) over time following randomisation (days) between continuation and discontinuation of antipsychotic medication in people with schizophrenia ( $k = 12$  trials,  $n = 2656$  participants). Vertical dashed lines mark the boundaries of time intervals over which effect estimates were made.

## References

- Baldessarini, R.J., Viguera, A.C., Faedda, G.L., Garver, D.L., Suppes, T., Tondo, L., Kammen, D. van, Wyatt, R.J., Gardner, D.M., 1995. Neuroleptic Withdrawal in Schizophrenic Patients. *Arch Gen Psychiatry* 52, 189–192.
- Buchanan, R.W., Kreyenbuhl, J., Kelly, D.L., Noel, J.M., Boggs, D.L., Fischer, B.A., Himelhoch, S., Fang, B., Peterson, E., Aquino, P.R., Keller, W., Schizophrenia Patient Outcomes Research Team (PORT), 2010. The 2009 schizophrenia PORT psychopharmacological treatment recommendations and summary statements. *Schizophr Bull* 36, 71–93.
- Chouinard, G., Samaha, A.-N., Chouinard, V.-A., Peretti, C.-S., Kanahara, N., Takase, M., Iyo, M., 2017. Antipsychotic-Induced Dopamine Supersensitivity Psychosis: Pharmacology, Criteria, and Therapy. *Psychotherapy and Psychosomatics* 86, 189–219.
- Correll, C.U., Rubio, J.M., Kane, J.M., 2018. What is the risk-benefit ratio of long-term antipsychotic treatment in people with schizophrenia? *World Psychiatry* 17, 149–160.
- Hasan, A., Falkai, P., Wobrock, T., Lieberman, J., Glenthøj, B., Gattaz, W.F., Thibaut, F., Möller, H.-J., WFSBP Task force on Treatment Guidelines for Schizophrenia, 2013. World Federation of Societies of Biological Psychiatry (WFSBP) guidelines for biological treatment of schizophrenia, part 2: Update 2012 on the long-term treatment of schizophrenia and management of antipsychotic-induced side effects. *The World Journal of Biological Psychiatry* 14, 2–44.



Leucht, S., Tardy, M., Komossa, K., Heres, S., Kissling, W., Salanti, G., Davis, J.M., 2012. Antipsychotic drugs versus placebo for relapse prevention in schizophrenia: A systematic review and meta-analysis. *Lancet* 379, 2063–2071.

Murray, R.M., Quattrone, D., Natesan, S., van Os, J., Nordentoft, M., Howes, O., Di Forti, M., Taylor, D., 2016. Should psychiatrists be more cautious about the long-term prophylactic use of antipsychotics? *Br J Psychiatry* 209, 361–365.

Parmar, M.K.B., Torri, V., Stewart, L., 1998. Extracting summary statistics to perform meta-analyses of the published literature for survival endpoints. *Stat Med* 17, 2815–2834.

Suppes, T., Baldessarini, R.J., Faedda, G.L., Tohen, M., 1991. Risk of recurrence following discontinuation of lithium treatment in bipolar disorder. *Arch Gen Psychiatry* 48, 1082–1088.

Tierney, J.F., Stewart, L.A., Gherzi, D., Burdett, S., Sydes, M.R., 2007. Practical methods for incorporating summary time-to-event data into meta-analysis. *Trials* 8, 16.